

<https://helda.helsinki.fi>

Changes in the latitudinal diversity gradient during the Great Ordovician Biodiversification Event

Kröger, Björn

2018-02

Kröger , B 2018 , ' Changes in the latitudinal diversity gradient during the Great Ordovician Biodiversification Event ' , Geology , vol. 46 , no. 2 , pp. 127-130 . <https://doi.org/10.1130/G39587.1>

<http://hdl.handle.net/10138/308336>

<https://doi.org/10.1130/G39587.1>

cc_by

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

1 Changes in the Latitudinal Diversity Gradient during the 2 Great Ordovician Biodiversification Event

3 **Björn Kröger***

4 *Finnish Museum of Natural History, PO Box 44, 00014 Helsinki, Finland*

5 *E-mail: bjorn.kroger@helsinki.fi

6 **ABSTRACT**

7 Near equatorial peak diversities are a prominent first-order feature of today's
8 latitudinal diversity gradient (LDG), but were not a persistent pattern throughout
9 geological time. In an analysis of Ordovician (485–445 Ma) fossil occurrences, an
10 equator-ward shift of the latitudinal diversity peak can be detected. A modern-type LDG
11 and out-of-the-tropics range shift pattern were synchronously established during
12 emerging icehouse conditions at the climax of the Great Ordovician Biodiversity Event
13 (GOBE). The changes in the LDG pattern and range shift trends can be best explained as
14 a consequence of global cooling during the Middle Ordovician and of diversification in
15 the tropical realm following a greenhouse period with temperatures too hot to support
16 diverse tropical marine life. These results substantiate a fundamental role of temperature
17 changes in establishing global first order diversity patterns.

18 **INTRODUCTION**

19 Life on earth today is most diverse at or near the equator and least diverse near the
20 poles (Pianka, 1966; Currie, 1991; Hillebrand, 2004; Mittelbach et al., 2007; Jablonski et
21 al. 2016). This latitudinal diversity gradient (LDG) is apparent in most groups of marine
22 and terrestrial organisms, but was not a persistent pattern throughout geological time

(e.g., Mannion et al., 2014). Strictly tropical peak diversities might be a feature of icehouse climate regimes, whereas warmer greenhouse regimes show generally shallow LDGs (e.g., Crame, 2002; Marcot et al., 2016; Jablonski et al., 2016)

These expectations can be tested against the fossil record from a time period during which one of the most dramatic and sustained Phanerozoic diversifications, the Great Ordovician Biodiversification Event (GOBE), took place. The GOBE was a ca. 40 Ma diversification interval during the Ordovician Period (485–444 m.y. ago) with a massive expansion in diversity at lower taxonomic ranks (species, genus, family level) (Fig. 1) and profound paleoecological changes within the marine biota (Servais et al., 2010). The diversification occurred against a background of changes in the physicochemical environment. In particular, it has been suggested that global cooling was one of the major driving factors, because this brought temperatures to levels more compatible with metazoan metabolism and skeletal growth (e.g., Trotter et al., 2008; Rasmussen et al., 2016). Hence, before cooling, large parts of the Ordovician seas, especially in the lower latitudes, would have been too hot to sustain highly diverse metazoan communities such as metazoan reefs (Webby, 2002). The GOBE, in this scenario, would represent a predominantly tropical diversification.

Ordovician diversity dynamics were previously analyzed mainly on a local, paleocontinent, or global scale (see e.g., Webby et al. 2004; Servais and Harper, 2013; Trubovitz and Stigall, 2016) and changes in LDG patterns can only indirectly be inferred from these data. Here, for the first time Ordovician marine diversity is analyzed across paleolatitudinal zones based on Cambrian–Silurian fossil occurrences documented in the Paleobiology Database (PaleoBioDB, <https://paleobiodb.org>). I reconstruct Ordovician

LDGs and their changes and ask whether the observed trends can be explained by changes in global temperatures, paleogeographical shifts, or changes in benthic habitats.

METHODS

The analysis was based on a download of the complete set of global, genus-level, Ordovician fossil occurrences from the Paleobiology Database on 26 November 2016. All accepted genus name entries in the PaleoBioDB were included in the analysis, without any additional filters set. Additionally, the complete set of 4081 genera from 773 fossil collections from the Cambrian Trempealeuan Regional Stage and the Silurian Rhuddanian Stage were downloaded. The stratigraphic data associated with each of the fossil occurrences were matched with stage level binned stratigraphic units in the RNames Database (<http://rnames.luomus.fi/>), resulting in 90,471 latest Cambrian to Rhuddanian occurrences (for time binning see Kröger and Lintulaakso, 2017). The genus occurrences are analyzed at stage- and stage-slice-level (Bergström et al., 2009) time resolution (average length 6.2 m.y. and 2.2 m.y., respectively).

PaleoBioDB collections are assigned to lithological and environmental classes. Based on these data I further classified all collections with sufficient lithology and environmental information into carbonate/siliciclastic and shallow/deep depositional environments (total = 13085 and 5739, respectively) and calculated their relative abundance (see the GSA Data Repository¹). Additionally, I used the PaleoBioDB data on lithology and environment for calculation of environmental heterogeneity using the HRel statistics of Wilcox (1973), (see the Data Repository).

All PaleoBioDB collections are explicitly attributed to paleoplates and are linked to paleogeographic latitudinal/longitudinal coordinates based on GPlates

(<https://www.gplates.org/>) rotations (Wright et al., 2013). The collections were binned into 15° and 30° paleolatitudinal zones without distinguishing between northern and southern paleo-hemispheres, because in most cases sample size for the northern paleo-hemisphere was not sufficient for analysis.

I estimated genus range shifts across paleolatitudinal zones and distinguished between “active” and “tectonic” shifts. The observed shift of a genus from one paleolatitudinal zone into another can be either a result of a tectonic shift of the genus location through time or can be caused by a active migrations or a range expansion of the genus, or both. Active shifts can be neutralized or reinforced by tectonic shifts of occurrences.

For distinction between tectonic and active range shift, I binned all genus occurrences into one-degree latitude/longitude quadrants and tracked their paleogeographical shift. Occurrences with identical quadrants in one stage (t_i) and its previous stage (t_{i-1}) and with different paleolatitudinal zones are interpreted as tectonically shifted. Occurrences with different latitude/longitude quadrants and different paleolatitudinal zones are interpreted as active range shifts.

Tectonic and active range shifts from one paleolatitudinal zone to another for times t_{i-1} to t_i , were counted with a unit length of ± 1 depending on the direction, such that positive values indicate pole-ward directions and negative values indicate equator-ward directions. In order to eliminate boundary effects, I compared the distance from the mean value of each latitudinal zone.

The diversities were estimated using Shareholder Quorum Subsampling (Alroy, 2010) (herein D_{sq}), Shannon Entropy Hill number (Chao et al., 2014) (herein D_{Chao}), and

the capture-mark-recapture (CMR) method (Liow and Nichols, 2010) (herein D_{CMR}).

These three methods were chosen because they represent independent approaches to account for sampling and preservation bias, which in their entirety give an estimate on the robustness of the estimated diversity pattern (Liow and Nichols, 2010; Chao et al. 2014, see the Data Repository). Because all three diversity measures, D_{SQS}, D_{Chao}, and D_{CMR}, exhibit a robust trend (Fig. DR1 in the Data Repository), only the D_{CMR} trends are described in detail in the main article, which additionally produced the most realistic (= largest) 95% confidence intervals. The downloaded data from the PaleoBioDB and the R code of the analysis are available at <https://doi.org/10.5281/zenodo.197057>.

RESULTS: CHANGING LDGs DURING THE COURSE OF THE DIVERSIFICATION

Ordovician diversity exhibits a trend with relatively low estimates for the first ~18 m.y. (Tremadocian–Dapingian), a steep increase during the remaining Middle Ordovician (Darriwilian), a Late Ordovician plateau, and a drastic decrease during the terminal Ordovician (Fig. 1). When partitioned into four 15° paleolatitudinal zones it becomes apparent that the bulk of the Ordovician diversity increase took place within the tropical realm (0–30°, Fig. 2). In temperate latitudes the GOBE is hardly visible.

This predominance of the tropical diversification can be interpreted as real because it is apparent in all diversity estimates explored herein (Fig. DR1), and it does not reflect changes in the relative abundance of carbonate/siliciclastic and shallow/deep depositional environments, in environmental and lithologic heterogeneity, or in paleogeographical shift (Fig. 3; Fig. DR5). However, although the pattern is not restricted

114 to a specific organismal clade because it is apparent in e.g., brachiopods, mollusks, and
115 trilobites, it is less pronounced or absent in planktic graptolites (Figs DR2 and DR3).

116 When comparing the per-time-bin diversity estimates against their paleolatitudinal
117 zone, an Ordovician LDG trend becomes apparent, the most important features of which
118 are an Early to Middle Ordovician (Tremadocian–Darriwilian) peak at the 15–45° zone, a
119 Late Ordovician (Sandbian–Katian) change toward higher tropical diversities with a peak
120 at 0–30°, and again a 15–30°-zone peak during the terminal Ordovician (Hirnantian) (Fig.
121 2; Fig. DR4). Notably, the Darriwilian marks not only the climax of the diversification,
122 but also a major pattern shift in Ordovician LDGs toward maximum values in the tropics.

123 LDG trends are an effect of changes in evolutionary rates and dispersal (see
124 Jablonski et al. 2016). Dispersal dynamics are driven by differential rates of active and
125 tectonic range shifts (see Methods). These dispersal factors are not correlated in the
126 Ordovician data, and tectonic shift appears to play a minor role in the overall picture (Fig.
127 3). The relative range shift pattern exhibits considerable variation during the Ordovician
128 with different trends near the equator and in temperate paleolatitudes (Fig. 2). In the high
129 tropics (0–15°), pole-ward shifts were relatively rare during the Early Ordovician but
130 became abruptly more common during the late Middle and Late Ordovician. In contrast,
131 in low tropical and temperate paleolatitudes, the trend exhibits initial (Tremadocian) high
132 rates of pole-ward shifts, relatively high rates of equator-ward shifts during the Floian-
133 Dapingian and an abrupt change toward moderate rates of pole-ward and equator-ward
134 shifts during the Darriwilian.

135 These trends reveal an intricate change in the general migration dynamics of the
136 Ordovician world with shifting diversity sinks and sources (Fig. 4). One prominent

feature of this pattern is the change of the high tropics from a sink during the Early Ordovician toward a source during the latter part of that period. This pattern is consistent with a change toward a modern-type “out of the tropics” scenario (Jablonski et al., 2006) during the Darriwilian concurrent with the main diversification pulse of the GOBE.

DISCUSSION: CHANGING LDGs AND GLOBAL COOLING

Previous analysis of Ordovician diversity trends below global level have focused on paleocontinents or paleo-oceanic regions that range across paleolatitudinal zones and often revealed heterogenous diversity pattern that are difficult to interpret (Miller, 1997; Webby et al., 2004). Only recently have synchronous Darriwilian diversification peaks within low paleolatitude regions of Baltica and Laurentia been detected in brachiopods; these results were used to infer global drivers, such as climatic cooling, intensified sea-water circulation, greater oxygenation and increased primary production (Trubovitz and Stigall, 2016). Here, I show that, although the Darriwilian peak diversification affected all latitudinal zones, it had by far the most profound impact and highest amplitude within the high tropics (Fig. 2). Combined with escalating rates of ex-tropical range shifts during the Middle Ordovician, this tropical diversification appears to be primarily caused by in situ origination, and in turn this tropical in situ origination must have been the quantitatively most important driver of the GOBE.

If this pattern is realistic, then tropical in situ origination is crucial to understanding the GOBE and hence the question of how this high tropical diversity was produced comes into focus. In a recent study and review this question was addressed with respect to the GOBE in general with oscillations between events of immigration and geographic differentiation suggested as one effective “speciation dynamo” (Stigall,

2017). Therein, geographic differentiation, conceptualized as tectonically induced regional isolation (e.g., Harper, 2010) and habitat heterogeneity (e.g., Miller and Mao, 1995) were considered to be key factors in promoting origination (Stigall, 2017). However, the habitat heterogeneity trends estimated herein are not related to the diversity pattern and generally the heterogeneity is not higher in the tropics (Fig. DR5). Because it is currently not known if regional differentiation was more intense in the tropics during the GOBE, these two factors, although not in conflict, cannot explain the LDG pattern detected herein.

Alternatively, geographic differentiation can be more generally conceptualized as ecological niche differentiation, including biologically-induced niche differentiation (*sensu* Sepkoski, 1988). With this more general conceptualization it is possible to set the GOBE into context with the temperature and evolutionary speed (TES) hypotheses of e.g., Allen et al. (2002), Clarke and Gaston (2006), and Brown (2014), which link ecological and evolutionary rates with temperature and predict highest diversities in the tropics. Generally, the Ordovician diversity trend is contrary to naive expectations from TES hypotheses, because they would predict decreasing ecological and evolutionary rates, and diversities and flattened LDG amplitudes under climatic cooling (Tittensor et al., 2010; Brown, 2014; Tittensor and Worm, 2016). This mismatch, however, can be best explained by the existing range limits of the published diversity-temperature dependencies. All published TES hypotheses are based on empirically observed positive diversity-temperature dependencies. These dependencies are constrained by Recent sea surface temperatures with a maximum of ~30 °C (e.g., Brown, 2014). Estimated Early Ordovician tropical sea surface temperatures are well beyond today's maximum values

(Trotter et al., 2008, Veizer and Prokoph, 2015). It is not known what diversity-temperature dependencies would be like in a hypothetical expanded temperature range, but thermal tolerances of complex multicellular organisms are limited, and today an upper limit for metazoans exists at ~45–47° (Ravaux et al., 2013). These limitations suggest that past diversity-temperature curves beyond modern temperature ranges had a parabolic shape with a diversity maximum in moderate temperatures and decreasing trajectories toward extreme temperatures. Past LDG's under hotter-than-today global regimes theoretically should reflect these expected trajectories with diversity peaks in temperate latitudes rather than near the equator. Data from Mesozoic and Cenozoic greenhouse intervals support this assumption (e.g., Markwick, 1998; Marcot et al., 2016) and the new data from the Ordovician are consistent with earlier assessments that tropical peak diversities are an exclusive feature of LDGs of icehouse periods (Mannion et al., 2014). Additionally, the Ordovician shift of peak LDG values toward the equator coincides with rapid initial expansion and diversification of metazoan reefs after a long period of dominance of reefs built by calcimicrobes, suggesting a (re)colonization of the low latitudes after a prolonged period of too-hot temperatures in the tropics (Webby, 2002; Kröger et al. 2017).

Hence, the mid-Ordovician cooling must have been a major factor for the diversification of the tropics. But why did this tropical diversification and LDG shift have such an overwhelming effect on global net diversity? An answer to this question is not easily found with the data at hand. The tropical mid-Ordovician diversification took place in a larger area and in regions with less intense seasonality compared with the temperate locus of the diversification during the Early Ordovician. The effects of these differences

on global Ordovician diversity are currently unknown and offer great potential for future investigations. Furthermore, the tropical mid-Ordovician diversification did not affect all clades. Brachiopods, mollusks, and trilobites diversified mainly in low latitudes and during the Darriwilian, whereas the tropical diversification of graptolites occurred earlier and the Darriwilian pulse is visible only in temperate latitudes. This asynchronicity between clades was known before (e.g., Webby et al., 2004) and new studies potentially will be key to answering this question in the future.

ACKNOWLEDGMENTS

I am indebted for the support by the Deutsche Forschungsgemeinschaft (grant KR, 2095/7–1). I am grateful for the critical reading of an earlier version of this manuscript and encouragement to implement CMR modeling for the diversity estimation by Lee Hsiang Liow (Oslo, Norway) and Melanie Hopkins (New York, USA). Jostein Starrfelt (Oslo, Norway) helped with the implementation, evaluation, and final exclusion of the “True Richness estimated using a Poisson Sampling model” (TRiPS) method during an earlier approach to calculate Ordovician diversities. Kari Lintulaakso (Helsinki, Finland) helped with the construction and realization of the RNames Database. This paper is a contribution to the IGCP 653 project.

REFERENCES CITED

- Allen, A.P., Brown, J.H., and Gillooly, J.F., 2002, Global biodiversity, biochemical kinetics, and the energetic-equivalence rule: *Science*, v. 297, p. 1545, <https://doi.org/10.1126/science.1072380>.
- Alroy, J., 2010, The shifting balance of diversity among major marine animal groups: *Science*, v. 329, p. 1191–1194, <https://doi.org/10.1126/science.1189910>.

- 229 Bergström, S.M., Chen, X., Gutiérrez-Marco, J.C., and Dronov, A., 2009, The new
230 chronostratigraphic classification of the Ordovician System and its relation to major
231 regional series and stages and to $\delta^{13}\text{C}$ chemostratigraphy: *Lethaia*, v. 42, p. 97–107,
232 <https://doi.org/10.1111/j.1502-3931.2008.00136.x>.
- 233 Brown, J.H., 2014, Why are there so many species in the tropics: *Journal of*
234 *Biogeography*, v. 41, p. 8–22, <https://doi.org/10.1111/jbi.12228>.
- 235 Chao, A., Gotelli, N.J., Hsieh, T.C., Sander, E.L., Ma, K.H., Colwell, R.K., and Ellison,
236 A.M., 2014, Rarefaction and extrapolation with Hill numbers: A framework for
237 sampling and estimation in species diversity studies: *Ecological Society of America*,
238 v. 84, p. 45–67.
- 239 Clarke, A., and Gaston, K.J., 2006, Climate, energy and diversity: *Proceedings.*
240 *Biological Sciences*, v. 273, p. 2257, <https://doi.org/10.1098/rspb.2006.3545>.
- 241 Crame, A.J., 2002, Evolution of taxonomic diversity gradients in the marine realm: A
242 comparison of Late Jurassic and Recent bivalve faunas: *Paleobiology*, v. 28, p. 184–
243 207, [https://doi.org/10.1666/0094-8373\(2002\)028<0184:EOTDGI>2.0.CO;2](https://doi.org/10.1666/0094-8373(2002)028<0184:EOTDGI>2.0.CO;2).
- 244 Currie, D.J., 1991, Energy and large-scale patterns of animal- and plant-species richness:
245 *American Naturalist*, v. 137, p. 27–49, <https://doi.org/10.1086/285144>.
- 246 Harper, D.A.T., 2010, The Ordovician brachiopod radiation: Roles of alpha, beta and
247 gamma diversity: *Geological Society of America Special Papers*, v. 466, p. 69–83.
- 248 Hillebrand, H., 2004, On the generality of the Latitudinal Diversity Gradient: *American*
249 *Naturalist*, v. 163, p. 192–211, <https://doi.org/10.1086/381004>.

- 250 Jablonski, D., Roy, K., and Valentine, J.W., 2006, Out of the Tropics: Evolutionary
251 dynamics of the Latitudinal Diversity Gradient: *Science*, v. 314, p. 102,
252 <https://doi.org/10.1126/science.1130880>.
- 253 Jablonski, D., Huang, S., Roy, K., and Valentine, J.W., 2016, Shaping the Latitudinal
254 Diversity Gradient: New perspectives from a synthesis of paleobiology and
255 biogeography: *American Naturalist*, v. 189, p. 1–12, <https://doi.org/10.1086/689739>.
- 256 Kröger, B., Desrochers, A., and Ernst, A., 2017, The reengineering of reef habitats during
257 the Great Ordovician Biodiversification Event: *Palaios*, v. 32, p. 584–599,
258 <https://doi.org/10.2110/palo.2017.017>.
- 259 Kröger, B., and Lintulaakso, K., 2017, RNames, a stratigraphical database designed for
260 the statistical analysis of fossil occurrences—The Ordovician diversification as a
261 case study: *Palaeontologica Electronica*, v. 20.1.1T, p. 1–12.
- 262 Liow, L.H., and Nichols, J.D., 2010, Estimating rates and probabilities of origination and
263 extinction using taxonomic occurrence data: Capture-mark-recapture (CMR)
264 approaches, *in* Alroy, J., and Hunt, G., ed., *The Paleontological Society Short*
265 *Course*, October 30, 2010, p. 81–94.
- 266 Mannion, P.D., Upchurch, P., Benson, R.B.J., and Goswami, A., 2014, The Latitudinal
267 Biodiversity Gradient through deep time: *Trends in Ecology & Evolution*, v. 29,
268 p. 42–50, <https://doi.org/10.1016/j.tree.2013.09.012>.
- 269 Marcot, J.D., Fox, D.L., and Niebuhr, S.R., 2016, Late Cenozoic onset of the Latitudinal
270 Diversity Gradient of North American mammals: *Proceedings of the National*
271 *Academy of Sciences of the United States of America*, v. 113, p. 7189–7194,
272 <https://doi.org/10.1073/pnas.1524750113>.

- 273 Markwick, P.J., 1998, Crocodilian diversity in space and time: The role of climate in
274 paleoecology and its implication for understanding K/T extinctions: *Paleobiology*,
275 v. 24, p. 470–497, <https://doi.org/10.1017/S009483730002011X>.
- 276 Miller, A.I., 1997, Dissecting global diversity patterns: Examples from the Ordovician
277 Radiation: *Annual Review of Ecology Evolution and Systematics*, v. 28, p. 85–104,
278 <https://doi.org/10.1146/annurev.ecolsys.28.1.85>.
- 279 Miller, A.I., and Mao, S., 1995, Association of orogenic activity with the Ordovician
280 radiation of marine life: *Geology*, v. 23, p. 305–308, [https://doi.org/10.1130/0091-](https://doi.org/10.1130/0091-7613(1995)023<0305:AOOAWT>2.3.CO;2)
281 [7613\(1995\)023<0305:AOOAWT>2.3.CO;2](https://doi.org/10.1130/0091-7613(1995)023<0305:AOOAWT>2.3.CO;2).
- 282 Mittelbach, G.G., et al., 2007, Evolution and the latitudinal diversity gradient: Speciation,
283 extinction and biogeography: *Ecology Letters*, v. 10, p. 315–331,
284 <https://doi.org/10.1111/j.1461-0248.2007.01020.x>.
- 285 Pianka, E.R., 1966, Latitudinal gradients in species diversity: A review of concepts:
286 *American Naturalist*, v. 100, p. 33–46, <https://doi.org/10.1086/282398>.
- 287 Rasmussen, C.M.Ø., Ullmann, C.V., Jakobsen, K.G., Lindskog, A., Hansen, J., Hansen,
288 T., Eriksson, M.E., Dronov, A., Frei, R., Korte, C., Nielsen, A.T., and Harper,
289 D.A.T., 2016, Onset of main Phanerozoic marine radiation sparked by emerging Mid
290 Ordovician icehouse: *Scientific Reports*, v. 6, p. 18884,
291 <https://doi.org/10.1038/srep18884>.
- 292 Ravaux, J., Hamel, G., Zbinden, M., Tasiemski, A.A., Boutet, I., Léger, N., Tanguy, A.,
293 Jollivet, D., and Shillito, B., 2013, Thermal limit for Metazoan life in question: *In*
294 *vivo* heat tolerance of the Pompeii Worm: *PLoS One*, v. 8, p. e64074,
295 <https://doi.org/10.1371/journal.pone.0064074>.

- 296 Sepkoski, J.J., 1988, Alpha, beta, or gamma: Where does all the diversity go?:
297 Palaeobiology, v. 14, p. 221–234, <https://doi.org/10.1017/S0094837300011969>.
- 298 Servais, T., and Harper, D.A.T., 2013, Early Palaeozoic biogeography and
299 palaeogeography: Geological Society of London Memoirs, v. 38, p. 476.
- 300 Servais, T., Owen, A.W., Harper, D.A.T., Kröger, B., and Munnecke, A., 2010, The
301 Great Ordovician Biodiversification Event (GOBE): The palaeoecological
302 dimension: Palaeogeography, Palaeoclimatology, Palaeoecology, v. 294, p. 99–119,
303 <https://doi.org/10.1016/j.palaeo.2010.05.031>.
- 304 Stigall, A.L., 2017, How is biodiversity produced? Examining speciation processes
305 during the GOBE: Lethaia, <https://doi.org/10.1111/let.12232>.
- 306 Tittensor, D.P., Mora, C., Jetz, W., Lotze, H.K., Ricard, D., Berghe, E.V., and Worm, B.,
307 2010, Global patterns and predictors of marine biodiversity across taxa: Nature,
308 v. 466, p. 1098–1101, <https://doi.org/10.1038/nature09329>.
- 309 Tittensor, D.P., and Worm, B., 2016, A neutral-metabolic theory of latitudinal
310 biodiversity: Global Ecology and Biogeography, v. 25, p. 630–641,
311 <https://doi.org/10.1111/geb.12451>.
- 312 Trotter, J.A., Williams, I.S., Barnes, C.R., Lécuyer, C., and Nicoll, R.S., 2008, Did
313 cooling oceans trigger Ordovician biodiversification? Evidence from conodont
314 thermometry: Science, v. 321, p. 550–554, <https://doi.org/10.1126/science.1155814>.
- 315 Trubovitz, S., and Stigall, A.L., 2016, Synchronous diversification of Laurentian and
316 Baltic rhynchonelliform brachiopods: Implications for regional versus global triggers
317 of the Great Ordovician Biodiversification Event: Geology, v. 44, p. 743–746,
318 <https://doi.org/10.1130/G38083.1>.

- 319 Veizer, J., and Prokoph, A., 2015, Temperatures and oxygen isotopic composition of
320 Phanerozoic oceans: *Earth-Science Reviews*, v. 146, p. 92–104,
321 <https://doi.org/10.1016/j.earscirev.2015.03.008>.
- 322 Webby, B.D., 2002, Patterns of Ordovician reef development, *in* Kiessling, W., et al.,
323 eds., *Phanerozoic Reef Patterns*: SEPM (Society for Sedimentary Geology), p. 129–
324 179, <https://doi.org/10.2110/pec.02.72.0129>.
- 325 Webby, B.D., Elias, R.J., Young, G.A., Neumann, B.E.E., and Kaljo, D., 2004, Corals, *in*
326 Webby, B.D., et al., eds., *The Great Ordovician Biodiversification Event*: New York,
327 Columbia University Press, p. 124–146, <https://doi.org/10.7312/webb12678-016>.
- 328 Wilcox, A.R., 1973, Indices of qualitative variation and political measurement: *The*
329 *Western Political Quarterly*, v. 26, p. 325–343, <https://doi.org/10.2307/446831>.
- 330 Wright, N., Zahirovic, S., Müller, R.D., and Seton, M., 2013, Towards community-driven
331 paleogeographic reconstructions: Integrating open-access paleogeographic and
332 paleobiology data with plate tectonics: *Biogeosciences*, v. 10, p. 1529–1541,
333 <https://doi.org/10.5194/bg-10-1529-2013>.

FIGURE CAPTIONS




Figure 1. Global genus-level diversity trends for the Ordovician Period with stratigraphic resolution at stage level (squares) and stage slice level (circles, Bergström et al., 2009). The peak diversification of the Great Ordovician Biodiversification Event (GOBE) was during the Darriwilian Stage. Data downloaded from Paleobiology Database (PaleoBioDB, <https://paleobiodb.org>) on 26 November 2016. D_{CMR}—diversity estimate from capture-mark-recapture (CMR) modeling approach with 95% confidence intervals (shaded areas). Stages: Tr—Tremadocian; Fl—Floian; Dp—Dapingian; Dw—Darriwilian; Sb—Sandbian; Ka—Katian, *—Hirnantian.





Figure 3. Trends in relative abundance of Ordovician fossil occurrences downloaded from Paleobiology Database (PaleoBioDB, <https://paleobiodb.org>) on 26 November 2016 from different environments and lithologies (carb.—carbonatic; silic. —siliciclastic), and direction of active/tectonic shift within two paleolatitudinal zones (Northern and Southern Hemispheres combined). Note the differences between active/tectonic shift, environmental trends, and diversity trends (see Fig. 2).

366



367

368 Figure 4. Summary of global change during the Ordovician. Paleogeographic
369 reconstructions from BugPlates (<http://www.geodynamics.no/bugs>). Diversity estimates
370 and range shift dynamics simplified from Figure 2 (see also Fig. DR4 [see footnote 1]).
371 Estimates are combined for northern and southern hemisphere and are symmetrically
372 plotted across equator for simplicity. n_{gen} —number of genera.

373

374 GSA Data Repository item 2018xxx, xxxxxxxx, is available online at
375 <http://www.geosociety.org/datarepository/2018/>, or on request from
376 editing@geosociety.org.